

Deep Vein Thrombosis in Focus: Resource Guide

Completing the Course: Implementing and Adhering to Guideline-based VTE Prophylaxis Across the Continuum of Care

Anderson FA Jr, Spencer FA. Risk factors for venous thromboembolism. *Circulation*. 2003;107:19-116.

Anderson FA Jr, Zayaruzny M, Heit JA, Fidan D, Cohen AT. Estimated annual numbers of US acute-care hospital patients at risk for venous thromboembolism. *Am J Hematol*. 2007;82:777-782.

Bahl V, Hu HM, Henke PK, Wakefield TW, Campbell DA Jr, Caprini JA. A validation study of a retrospective venous thromboembolism risk scoring method. *Ann Surg*. 2009 (Epub ahead of print)

Bergqvist D, Agnelli G, Cohen AT, et al. Duration of prophylaxis against venous thromboembolism with enoxaparin after surgery for cancer. *N Engl J Med*. 2002;346(13):975-980.

Caprini JA. Thrombosis risk assessment as a guide to quality patient care. *Dis Mon*. 2005;51:70-78.

Cohen AT, Tapson VF, Bergmann JF, et al. Venous thromboembolism risk and prophylaxis in the acute hospital care setting (ENDORSE study): a multinational cross-sectional study. *Lancet*. 2008;371(9610):387-394.

Eikelboom JW, Quinlan DJ, Douketis JD. Extended-duration prophylaxis against venous thromboembolism after total hip or knee replacement: a meta-analysis of the randomised trials. *Lancet*. 2001;358:9-15.

Geerts WH, Bergqvist D, Pineo GF, et al. Prevention of venous thromboembolism: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. (8th Edition). *Chest*. 2008;133(6 suppl):381S-453S.

Heit JA, Silverstein MD, Mohr DN, Petterson TM, O'Fallon M, Melton LJ III. Risk factors for deep vein thrombosis and pulmonary embolism: A population-based case-control study. *Arch Intern Med*. 2000;160:809-815.

Hull RD, Schellong SM, Tapson VF, et al. Presented at: XXist Congress of the International Society on Thrombosis and Haemostasis; July 6-12, 2007; Geneva, Switzerland. Abstract O-S-001.

Kucher N, Koo S, Quiroz R, et al. Electronic alerts to prevent venous thromboembolism among hospitalized patients. *N Engl J Med*. 2005;352(10):969-977.

Piazza G, Rosebaum EJ, Pendergast W, et al. Physician alerts to prevent symptomatic venous thromboembolism in hospitalized patients. *Circulation*. 2009;119(16):2196-2201.

Spencer FA, Lessard D, Emery C, Reed G, Goldberg RJ. Venous thromboembolism in the outpatient setting. *Arch Intern Med*. 2007;167(14):1471-1475.

Tapson VF, Decousus H, Pini M, et al. Venous thromboembolism prophylaxis in acutely ill hospitalized medical patients: findings from the International Medical Prevention Registry on Venous Thromboembolism. *Chest*. 2007;132(3):936-945.

Tooher R, Middleton P, Pham C, et al. A systematic review of strategies to improve prophylaxis for venous thromboembolism in hospitals. *Ann Surg*. 2005;241:397-415.

US Department of Health and Human Services. The Surgeon General's Call to Action to Prevent Deep Vein Thrombosis and Pulmonary Embolism 2008.
<http://www.surgeongeneral.gov/topics/deepvein/calltoaction/call-to-action-on-dvt-2008.pdf>.
Accessed November 12, 2009.

Yu HT, Dylan ML, Lin J, Dubois RW. Hospitals' compliance with prophylaxis guidelines for venous thromboembolism. *Am J Health Syst Pharm*. 2007;64:69-76.